# **KernMedical**

#### Introduction

This case presents an image of a condition that is rare and particularly severe, as shown by free air not only in the right renal parenchyma, but also extending outside the capsule, around the renal vasculature, and into the left perirenal space.

#### Objectives

A 58-year- old male presented to an outside hospital with altered mental status and right flank pain for three days. Septic work up, including computed tomography of the abdomen and pelvis, were significant for diabetic ketoacidosis, pyelonephritis, and significant air replacing much of the right kidney, consistent with emphysematous pyelonephritis (Image). The patient was transferred to our facility for a higher level of care. The patient was stabilized, given intravenous (IV) antibiotics, and admitted to the intensive care unit with a diagnosis of septic shock secondary to emphysematous pyelonephritis.

## Massive Emphysematous Pyelonephritis

Adria Ottoboni<sup>1</sup>, MD; Shannon Anderson<sup>2</sup>, MD; Natalie Peña-Brockett<sup>3</sup>, MS; Phillip Aguìñiga-Navarrete<sup>3</sup> <sup>1</sup> Faculty, Emergency Medicine; <sup>2</sup> PGY IV, Emergency Medicine; <sup>3</sup> Research Assistant, Emergency Medicine Research Assistant Program

#### Imaging

Image 1: Coronal view of a CT of the abdomen and pelvis in the lung window showing bilateral emphysema



**Image 2**: Axial view of Ct of the abdomen and pelvis without contrast showing emphysema replacing the right kidney.



### Conclusions

Emphysematous pyelonephritis is a relatively rare infection, seen only 1-2 times per year in a typical busy urological department in the United States. It affects patients with diabetes in 95% of cases. E. coli and Klebsiella account for over 90% of cases, although Proteus mirabilis, Pseudomonas, and Streptococcus are also seen. Gas accumulates due to rapid necrosis of the renal parenchyma and perirenal tissue, as opposed to gas appearing as a byproduct of anaerobic bacteria as is the case in necrotizing fasciitis. The condition is fatal if not treated appropriately, and the mainstay of treatment is nephrectomy in conjunction with IV antibiotics for severe, disseminated infection.

Tal May et al. Carmel Medical Center, Department of Urology, 7 Michal Street, Haifa, Israelb Carmel Medical Center, Department of Radiology, 7 Michal Street, Haifa, Israel Demonstrative Imaging of Emphysematous Cystitis. Urology Case Reports 6 (2016) 56e57

1988

Huang J, Tseng C. Emphysematous pyelonephritis: clinicoradiological classification, management, prognosis, and pathogenesis. Arch Intern Med. 2000;160:797-805.

#### References

Lyne M Ouellet, MD, Michael P Brook, MD. Emphysematous Pyelonephritis: An Emergency Indication for the Plain Abdominal Radiograph. Annals of Emergency Medicine 17:7 July